

Javier Carvajal-Rojas¹, Anne Fayolle²
 CIMAT¹, University of Utah²

Abstract: I introduce a notion of tame ramification for general finite covers. When specialized to the separable case, it extends to higher dimensions the classical notion of tame ramification for Dedekind domains and curves. However, when applied to the Frobenius map, it naturally yields the notion of centers of F -purity, which lets us describe how these behave under finite covers. This is joint work with Javier Carvajal-Rojas

Theorem: R and S Dedekind domains, $\mathfrak{T} = \text{Tr}$ the trace.
 $\left\{ \begin{array}{l} S/R \text{ is tamely} \\ \mathfrak{T}\text{-ramified over } \mathfrak{p} \end{array} \right\} \iff \left\{ \begin{array}{l} S/R \text{ is tamely} \\ \text{ramified over } \mathfrak{p} \end{array} \right\}$

Definition: R of characteristic $p > 0$, $\mathfrak{a} \subset R$ an ideal, and $\phi \in \text{Hom}(F_*^e R, R)$ surjective. We say \mathfrak{a} is ϕ -compatible if $\phi(F_*^e \mathfrak{a}) \subset \mathfrak{a}$.

Notation and assumptions:

- $\theta: R \rightarrow S$ is a finite ring extension and \mathfrak{T} is in $\text{Hom}_R(S, R)$.
- If R is a ring of characteristic $p > 0$ then $F: R \rightarrow R$ is the Frobenius morphism, F^e its e^{th} iteration.
- $F_*^e R$ is an R -module which is R as a set but R acts on it via F^e as follows: $rF_*^e r' = F_*^e r^{p^e} r'$.
- All rings are Noetherian and those of char. $p > 0$ are F -finite i.e. $F_*^e R$ is a finite R module for all e .

Definition: Let $\mathfrak{a} \subset R$, then

$$\mathfrak{a}^{\mathfrak{T}} := \{s \in S \mid \mathfrak{T}(sS) \subset \mathfrak{a}\} \subset S$$

is the largest ideal of S whose image under \mathfrak{T} is in \mathfrak{a} .

Proposition:

- $\mathfrak{a}^{\mathfrak{T}}$ is a proper ideal if and only if $\mathfrak{T}(S) \not\subseteq \mathfrak{a}$.
- If $\mathfrak{p} \in \text{Spec } R$ and $\mathfrak{T}(S) \not\subseteq \mathfrak{p}$ then $\mathfrak{p}^{\mathfrak{T}} \cap R = \mathfrak{p}$. Moreover if $\mathfrak{q} \in \text{Ass}(S/\mathfrak{p}^{\mathfrak{T}})$ then $\mathfrak{q} \cap R = \mathfrak{p}$.

Definition: S/R is tamely \mathfrak{T} -ramified over $\mathfrak{p} \in \text{Spec } R$ if $\mathfrak{p}^{\mathfrak{T}} = \sqrt{\mathfrak{p}S}$.

Facts:

- (R, ϕ) is F -regular $\iff (0)$ and (1) are the only ϕ -compatible ideals.
- The set of ϕ -compatible ideals tells us where the singularities of (R, ϕ) are the most severe.
- These ideals are radical and closed under taking minimal primes \implies can focus on the prime ones, the centers of F -purity (CFPs).
- $\mathfrak{a} \subset R$ is ϕ -compatible $\iff \mathfrak{a}^\phi = \mathfrak{a}$ in $F_*^e R$.

Definition [3]: For $\mathfrak{p} \in \text{Spec } R$ a CFP, $\tau_{\mathfrak{p}}(R, \phi)$ is the smallest ϕ -compatible ideal \mathfrak{a} of R such that $\mathfrak{a} \not\subseteq \mathfrak{p}$.

Definition [2]: A transposition of pairs $(\theta, \mathfrak{T}): (R, \phi) \rightarrow (S, \psi)$ is a commutative diagram

$$\begin{array}{ccc} F_*^e S & \xrightarrow{\psi} & S \\ F_*^e \mathfrak{T} \downarrow & & \downarrow \mathfrak{T} \\ F_*^e R & \xrightarrow{\phi} & R \end{array}$$

with (R, ϕ) as above, $\text{char } S = p$, and ψ surjective.

Remark: For $\mathfrak{p} \in \text{Spec } R$, following the previous diagram from the bottom right gives

$$\begin{array}{ccc} (F_*^e \mathfrak{p}^\phi)^{F_*^e \mathfrak{T}} = F_*^e (\mathfrak{p}^{\mathfrak{T}})^\psi & \xrightarrow{\psi} & \mathfrak{p}^{\mathfrak{T}} \\ F_*^e \mathfrak{T} \downarrow & & \downarrow \mathfrak{T} \\ F_*^e \mathfrak{p}^\phi & \xrightarrow{\phi} & \mathfrak{p} \end{array}$$

when $\mathfrak{p} = \mathfrak{p}^\phi$, $F_*^e \mathfrak{p}^{F_*^e \mathfrak{T}} = F_*^e (\mathfrak{p}^{\mathfrak{T}})^\psi$ so $\mathfrak{p}^{\mathfrak{T}} = (\mathfrak{p}^{\mathfrak{T}})^\psi \implies \mathfrak{p}^{\mathfrak{T}}$ is ψ -compatible.

Theorem: $(\theta, \mathfrak{T}): (R, \phi) \rightarrow (S, \psi)$ a transposition of pairs, \mathfrak{p} a CFP for (R, ϕ) , TFAE:

- All $\mathfrak{q} \in \text{Spec } S$ lying above \mathfrak{p} are CFPs for (S, ψ) and $\mathfrak{T}(\tau_{\mathfrak{q}}(S, \psi)) = \tau_{\mathfrak{p}}(R, \phi)$ for all such \mathfrak{q} .
- S/R is tamely \mathfrak{T} -ramified over \mathfrak{p}

Remark: The theorem generalizes to the case ϕ, ψ are not surjective by updating the definition of CFPs to $\mathfrak{p} \in \text{Spec } R$ is a CFP if $\phi(F_*^e \mathfrak{p}) \subset \mathfrak{p}$ and $\phi_{\mathfrak{p}}$ is surjective.

References:

- J. Carvajal-Rojas and A. Fayolle: On tame ramification and centers of F -purity, arXiv e-prints (2023).
- K. Schwede and K. Tucker: On the behavior of test ideals under finite morphisms, J. Algebraic Geom. 23 (2014), no. 3, 399–443. 3205587
- D. Smolkin: A new subadditivity formula for test ideals, J. Pure Appl. Algebra 224 (2020), no. 3, 1132–1172.

Acknowledgements: The second named author is partially supported by an NSERC doctoral fellowship.